

## Emergent classicality in continuous quantum measurements

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2013 New J. Phys. 15 013005

(<http://iopscience.iop.org/1367-2630/15/1/013005>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 152.78.1.74

This content was downloaded on 14/10/2016 at 13:02

Please note that [terms and conditions apply](#).

You may also be interested in:

[Chaos in circuit QED: decoherence, localization, and nonclassicality](#)

Jonas Larson and Duncan H J O'Dell

[Bragg spectroscopic interferometer and quantum measurement-induced correlations in atomic Bose–Einstein condensates](#)

M D Lee, S Rist and J Ruostekoski

[Dynamical formation and interaction of bright solitary waves and solitons in the collapse of Bose–Einstein condensates with attractive interactions](#)

B J Dabrowska-Wüster, S Wüster and M J Davis

[Quantum optics with ultracold quantum gases: towards the full quantum regime of the light–matter interaction](#)

Igor B Mekhov and Helmut Ritsch

[Strong-coupling dynamics of Bose–Einstein condensate in a double-well trap](#)

V O Nesterenko, A N Novikov and E Suraud

[Quantum dynamics of atomic bright solitons under splitting and re-collision, and implications for interferometry](#)

A D Martin and J Ruostekoski

## Emergent classicality in continuous quantum measurements

Juha Javanainen<sup>1</sup> and Janne Ruostekoski<sup>2</sup>

<sup>1</sup> Department of Physics, University of Connecticut, Storrs, CT 06269-3046, USA

<sup>2</sup> School of Mathematics, University of Southampton, Southampton, SO17 1BJ, UK

E-mail: [jj@phys.uconn.edu](mailto:jj@phys.uconn.edu)

*New Journal of Physics* **15** (2013) 013005 (9pp)

Received 18 September 2012

Published 3 January 2013

Online at <http://www.njp.org/>


doi:10.1088/1367-2630/15/1/013005

**Abstract.** We develop a classical theoretical description for nonlinear many-body dynamics that incorporates the back-action of a continuous measurement process. The classical approach is compared with the exact quantum solution in an example with an atomic Bose–Einstein condensate in a double-well potential where the atom numbers in both potential wells are monitored by light scattering. In the classical description, the back-action of the measurements appears as diffusion of the relative phase of the condensates on each side of the trap. When the measurements are frequent enough to resolve the system dynamics, the system behaves classically. This happens even deep in the quantum regime, and demonstrates how classical physics emerges from quantum mechanics as a result of measurement back-action.

It is now possible to carry out experiments with systems composed of a small and controllable number of atoms, and eventually the interface between quantum mechanics and classical physics [1–3] must be confronted. Measurement back-action is an essential element of quantum mechanics, and may play a key role also in the transition from quantum mechanics to classical physics [4–7]. In this paper, we introduce a theoretical method to account for measurement back-action in nonlinear dynamics [8–13] classically, using a continuously monitored [14] Bose–Einstein condensate (BEC) in a double-well trap as an explicit example. When the



Content from this work may be used under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 licence](https://creativecommons.org/licenses/by-nc-sa/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



**Figure 1.** Left: BEC in a double-well potential with tunneling amplitude  $J$ , atom–atom interaction strength  $U$  and energy difference per atom between the wells  $\epsilon$  indicated. Right: the basic idea of detection of the atoms in one trap. Since the detection light is far-off resonance, the dominant effect of the atoms is to alter the phase of the light passing through the sample by an amount proportional to atom number. The phase shift is then detected interferometrically. The amplitude of the light passing to the photodetector is proportional to the number of atoms, so that the rate of photon counts is proportional to the square of the atom number. The scheme is heavily simplified. For instance, mode matching of the light to the condensate is not considered.

continuous measurements are strong enough that the system dynamics can be resolved, the classical description and a full quantum analysis agree well even deep within what might appear to be the domain of quantum mechanics.

We employ the minimal two-mode model [15] to describe the Josephson oscillations of the atoms between the potential wells [16–20], see figure 1. Given the hopping amplitude  $J$  between the wells, the strength of atom–atom interaction inside a well  $U$  and the imbalance in the energies of the atoms between the sides  $\epsilon$ , the Hamiltonian is

$$H = -J(a^\dagger b + b^\dagger a) + U(a^\dagger a^\dagger a a + b^\dagger b^\dagger b b) + \epsilon(a^\dagger a - b^\dagger b). \quad (1)$$

Here  $a$  and  $b$  are the boson operators for the atoms in the two wells, and we set  $\hbar = 1$ .

The atom numbers in the two wells are monitored by coherent scattering of off-resonant light; consult figure 1. Counting of photons on two detectors, one for each potential well, constitutes the final measurement. In our quantum description, we consider the atoms as an open system and model the back-action of the measurements by adding a linear relaxation term to the equation of motion of the density operator of the atoms,

$$\dot{\rho} = -i[H, \rho] + \mathcal{L}\rho. \quad (2)$$

The most general relaxation term  $\mathcal{L}\rho$  such that a density operator remains valid during the time evolution is of the Lindblad form [21, 22]

$$\mathcal{L}\rho = \sum_i \left[ L_i \rho L_i^\dagger - \frac{1}{2}(\rho L_i^\dagger L_i + L_i^\dagger L_i \rho) \right], \quad (3)$$

where  $L_i$  are operators of the double-well system. In the present model [14], we have the Lindblad operators  $L_a = \sqrt{\Gamma} a^\dagger a$  and  $L_b = \sqrt{\Gamma} b^\dagger b$ , indicating that the scattering rates of photons from the two wells are  $\Gamma \langle (a^\dagger a)^2 \rangle$  and  $\Gamma \langle (b^\dagger b)^2 \rangle$ , respectively. If the light is predominantly scattered into a narrow cone in the forward direction,  $\Gamma$  also specifies the strength of the measurements. It can be adjusted, for instance, by varying the intensity or the frequency of the probing light.

One could develop a parallel classical description using various phase-space representations such as the Wigner function for the ordinary [22–24] or the SU(2) [25–27] coherent states, or the number-phase Wigner representation [28]. We choose the standard Wigner function because it is generally used in quantum optics and is not specifically tailored to the present system. We begin by rewriting the master equation (2) equivalently in terms of the Wigner function  $W(\alpha, \alpha^*, \beta, \beta^*; t)$  [22, 24], where  $\alpha$  and  $\beta$  are complex numbers corresponding to the operators  $a$  and  $b$ . In the equation of motion of the Wigner function,  $\alpha, \alpha^*$  and  $\beta, \beta^*$  behave like canonical pairs with Poisson brackets such as  $\{\alpha, \alpha^*\} = -i$ . We then change to four real variables  $N, \Phi, Z$  and  $\varphi$  defined by

$$\alpha = \sqrt{\frac{1}{2}N - Z} e^{i(\Phi - (1/2)\varphi)}, \quad \beta = \sqrt{\frac{1}{2}N + Z} e^{i(\Phi + (1/2)\varphi)}. \quad (4)$$

This transformation between the variables is canonical and results in Poisson brackets such as  $\{Z, \varphi\} = 1$ . To the extent possible, we regard the Wigner function as the classical joint probability density for the total atom number  $N$ , imbalance of atom numbers between the sides of the double well  $Z$  ( $Z \in [-N/2, N/2]$ ), overall phase of the two BECs  $\Phi$  and phase difference of the BECs between the two potential wells  $\varphi$ . The global phase  $\Phi$  may be dropped, as it has no dynamics, nor has it any relevance to the dynamics of the other variables. We define  $\chi = UN$ ,  $z = 2Z/N$  ( $z \in [-1, 1]$ ), and expand the equation of motion of the Wigner function up to and including the order  $\mathcal{O}(N^{-1})$  in the formally large atom number  $N$ . The equation of motion

$$\partial_t W = \left[ -\frac{\partial h}{\partial \varphi} \partial_z + \frac{\partial h}{\partial z} \partial_\varphi + \Gamma \partial_{\varphi\varphi} \right] W \quad (5)$$

follows for the remaining distribution function  $W(z, \varphi)$ , with the effective one-particle Hamiltonian

$$h(z, \varphi) = z^2 \chi - 2\epsilon z - 2J\sqrt{1 - z^2} \cos(\varphi). \quad (6)$$

Importantly, the quantum mechanical back-action of the measurements of the particle numbers is now accounted for by classical diffusion of the relative phase of the BECs proportional to  $\Gamma \partial_{\varphi\varphi}$ . This is, perhaps, as expected: the relevant measurement result is the difference of the photon counting rates on the two detectors, and this is proportional to the expectation value of the operator

$$\Delta \hat{N} = (a^\dagger a)^2 - (b^\dagger b)^2 = (a^\dagger a + b^\dagger b)(a^\dagger a - b^\dagger b). \quad (7)$$

But the first factor in the final form is the conserved total atom number, so the measured operator is actually the difference of atom numbers between the sides of the trap. Given the (in quantum mechanics, qualitative) conjugate relation between number and phase, measurement of atom number difference should indeed increase the uncertainty of the relative phase.

We compare the quantum and classical evolutions in the case when the system is first prepared to the ground state at zero temperature in the presence of an energy imbalance,

$\epsilon \neq 0$ , and the imbalance is suddenly removed by setting  $\epsilon = 0$ . The atoms then start oscillating between the two sides of the potential well.

In our quantum analysis the evolution of the density operator according to (2) is solved using quantum trajectory simulations [29–31]. For every run of the simulation we have a state vector that undergoes ‘quantum jumps’ at random times according to a precisely defined probabilistic law. As discussed in, say, [22, 32], here we also assume an ‘efficient’ measurement process in which every scattered photon gets detected. Then a quantum trajectory simulation with its quantum jumps produces a faithful simulation of an individual experimental run, each quantum jump corresponding to a detection of a photon. Even though the master equation (2) is not conditioned on the results of the measurements and as such describes an experiment in which the measurement results are discarded, the results are built into each quantum trajectory.

In a real experiment, one would record a stream of photon counts, and the population imbalance  $z(t)$  has to be inferred from them in some manner. For instance, if frequent photon counts merge into an essentially continuous photocurrent that varies on a time scale of the back-and-forth oscillations of the atoms, it would be sensible to regard the difference of the photocurrents coming from the two detectors as a measure of the instantaneous population imbalance. However, for the sake of a clean illustration we analyze our quantum simulations differently: given a quantum trajectory, a time-dependent state vector, we use the expectation value of  $\hat{z} = (a^\dagger a - b^\dagger b)/N$  to represent the value of the experimentally inferred population imbalance  $z(t)$ . Evidently, this will work the better the more numerous the quantum jumps are on the time scale of the variation of  $z(t)$ .

In the classical analysis, we take into account the quantum fluctuations in the initial state in the same way as is customary with the truncated Wigner approximation (TWA) [33–36]. Thus, suppose that equations (5) and (6) were written in terms of the variables  $Z$  and  $\varphi$  instead of  $z$  and  $\varphi$ . We expand the corresponding Hamiltonian  $\mathcal{H}(Z, \varphi)$  to second order in  $Z$  and  $\varphi$  around the classical lowest-energy state in the presence of the initial energy imbalance  $\epsilon \neq 0$  between the sides. Since the classical variables  $Z$  and  $\varphi$  are a canonical pair, we may requantize by postulating the commutator  $[Z, \varphi] = i$ .  $\mathcal{H}(Z, \varphi)$  thereby becomes the quantum Hamiltonian of a harmonic oscillator. The Wigner function corresponding to the ground state of this oscillator is used as the initial distribution of the variables  $Z$  and  $\varphi$  and leads to the initial condition for the distribution of the variables  $z$  and  $\varphi$

$$W_0(z, \varphi) = \frac{N}{2\pi} \exp \left[ -\frac{N}{2} \left( \frac{(z-f)^2}{\xi} + \xi \varphi^2 \right) \right]. \quad (8)$$


We take  $\chi > 0$  and choose  $\epsilon$  in such a way that the classical equilibrium has the phase  $\varphi = 0$  and a predetermined population imbalance  $z = f$ . The parameter

$$\xi = \frac{1-f^2}{\sqrt{1+(1-f^2)^{3/2}\chi/J}} \quad (9)$$

reflects the size of the ground state wave function of the underlying harmonic oscillator.

The evolution (5) with the initial Wigner function  $W_0(z, \varphi)$  is solved by unraveling the dynamics into stochastic trajectories [26, 33, 34, 37]. If there were no phase diffusion in (5), we would have an ensemble of trajectories  $(z(t), \varphi(t))$  evolving under the effective Hamiltonian  $h$ , which makes the usual TWA. Given the diffusion, there exists a precise mathematical correspondence between (5) and the stochastic differential equations

$$dz = 2J\sqrt{1-z^2} \sin \varphi dt, \quad (10)$$




**Figure 2.** A classical (upper trace, blue) and a quantum (lower trace, red) trajectory. The parameters are  $N = 100$ ,  $\chi/J = 10$ ,  $f = 0.5$  and  $\Gamma = 0.003$ .

$$d\varphi = -2z \left( \chi + J \frac{\cos \varphi}{\sqrt{1-z^2}} \right) dt + \sqrt{2\Gamma} dW(t), \quad (11)$$

where  $dW(t)$  denotes a random increment of the Wiener process over the interval  $dt$  [37]. The evolution of the distribution  $W(z, \varphi; t)$  from the initial condition  $W_0(z, \varphi)$  according to (5) is obtained from an ensemble of stochastic trajectories. Specifically, the initial state of each trajectory  $(z, \varphi)$  is sampled from the distribution  $W_0(z, \varphi)$ , the dynamics of the trajectory  $(z(t), \varphi(t))$  is determined from (10) and (11), and the expectation value of any quantity over the distribution  $W(z, \varphi; t)$  is correctly obtained by averaging the value of the quantity over a large number of such trajectories. The stochastic dynamics is integrated numerically using the Milstein algorithm [37].

We are now in a position to state precisely what ‘classical description’ means in this context: unlike quantum mechanics, the analysis conforms to classical logic. The usual mean-field theory [17] is a classical description, as are the TWA and our present approach. The initial Wigner function  $W_0$  is a valid classical probability distribution even when it approximately incorporates quantum fluctuations of the atomic populations in the initial state. Moreover, subsequent evolution according to (5) that also accounts for measurement back-action keeps the probability distribution function valid at all future times. Correspondingly, we have been able to represent the complete solution in terms of classical stochastic trajectories. Such a trajectory exists objectively even without observations, but otherwise a similar correspondence between simulations and experiments prevails in the classical description as in quantum mechanics: each individual stochastic realization would be a representative outcome of an individual experiment.

Figure 2 shows a classical (blue upper trace) and a quantum rendition (red lower trace) of a single experiment. In the mean-field theory, and for each individual TWA trajectory, one expects strictly periodic oscillations in the classical trace. The deviations from periodicity originate from the phase diffusion. On the other hand, the measurement strength  $\Gamma/J = 0.003$  is chosen such that each photon detector produces about ten clicks during a typical back-and-forth swing of the atoms, so that the quantum trace is a fair representation of a measurement record. We have deliberately picked these random examples so that qualitatively similar classical and quantum behaviors with occasional trapping of the populations [17] in one or the other well are seen.



**Figure 3.** Quantum (solid red lines) and classical (dashed blue lines) expectation values of the population imbalance  $z$  as a function of time  $t$ , all averaged over 1000 trajectories. The upper half of the figure is for atom number  $N = 100$  and the lower half for  $N = 10$ . Graphs (a) and (c) are without continuous monitoring,  $\Gamma/J = 0$ ; graphs (b) and (d) correspond to the observation strengths  $\Gamma/J = 0.003$  and  $0.3$ . The atom–atom interaction strength is  $\chi/J = 10$ , and the initial population imbalance is  $f = 0.5$ . The insets demonstrate the quality of the quantum–classical agreement in the collapse region of traces (b) and (d).

Nevertheless, even if we are able to model individual experiments both classically and quantum mechanically, these are stochastic stimulations. In order to construct expectation values of experimental observables for quantitative comparisons we have to average over many trajectories. In figure 3, we demonstrate simulation results for population imbalance when averaged over 1000 trajectories. The upper two panels are for  $N = 100$  atoms and the lower two are for  $N = 10$ , while the differences between graphs (a)–(d) are in the measurement strength  $\Gamma$ .

Panel (a) for  $N = 100$  atoms plots the quantum and classical expectation values of the population imbalance  $z(t)$  with no measurements,  $\Gamma = 0$ . The quantum result then represents the usual quantum average of the operator  $\hat{z} = (a^\dagger a - b^\dagger b)/N$ , and has nothing to do with the (nonexistent) continuous monitoring. The quantum and classical graphs both show a collapse up to  $Jt \simeq 10$ , but only the quantum expectation value has a revival at around  $Jt = 70$ . In panel (b), we turn up the measurements strength to  $\Gamma/J = 3 \times 10^{-3}$ . Both detectors then report about ten photons during a characteristic oscillation time of the atoms between the traps, so that the

oscillations of the atoms could be resolved by monitoring the light scattered off the atoms. The quantum expectation value of  $\hat{z}$  should therefore be a passable representation of the population imbalance inferred from photon counts. The collapse part, quantum and classical, is almost unchanged by the measurements, but the quantum revival has disappeared. The quantum and classical analyses now agree very well.

We finally move on to the atoms number  $N = 10$ . In panel (c), we show the population imbalances as a function of time for both the quantum theory and the classical theory without continuous monitoring,  $\Gamma = 0$ . There is little in common with the two traces, and one would conclude that the classical description is totally off. The possible reason is that with a decreasing atom number the quantum revivals start overlapping and produce irregular-looking behavior that is absent in the classical approach. It appears that we are deep in the domain of quantum mechanics. However, in panel (d), we have continuous measurements with the strength  $\Gamma/J = 0.3$ , which again gives about ten photon counts during a back-and-forth oscillation of the atoms; and again, the classical and quantum results agree.

We have surveyed the behavior of the system for parameter values comparable to those in figures 2 and 3. In the absence of measurements,  $\Gamma = 0$ , we have the usual TWA and the corresponding full quantum theory. The TWA relies on a stochastic description and cannot fully incorporate quantum effects. The generic observation is that TWA agrees well with the exact solution for short times when the stochastic field is highly occupied, but it becomes progressively less accurate a model of quantum dynamics as the average mode population becomes smaller and the simulation time scales longer [33–35]. There are also examples, however, especially in lower-dimensional systems in which case the TWA simulations have been successful in providing qualitative descriptions of experimental findings even in the strongly fluctuating quantum limit [38].

In figure 3, in traces (a) with  $N = 100$  and  $\Gamma = 0$  we find reasonable agreement between the TWA and quantum solutions up to the quantum revival around  $Jt = 70$ , while in figure 3(c) with  $N = 10$  the two cases part ways after one back-and-forth swing of the atoms at  $Jt \simeq 1$ , a time that is hard to discern in the figure at all. Secondly, as the measurement strength is increased from zero, the changes in the quantum results start becoming qualitatively visible at times when there have been, on average, a few counts on both detectors. The classical results also change with increasing  $\Gamma$ , but the changes are less conspicuous. For example, with  $N = 100$  the maximum amplitude of the quantum revival at  $Jt \simeq 70$  has decreased to half its value in figure 3(a) for  $\Gamma/J = 2 \times 10^{-5}$ , at which point the average time between photon counts at each detector is about  $20/J$ . Let us next consider  $\Gamma/J = 3 \times 10^{-3}$ , the value used in figure 3(b) for  $N = 100$ . Since our definition of the parameter  $\Gamma$  is such that the photon counting rate on the detectors is proportional to the square of the atom number, for  $N = 10$  this measurement strength is still modest, with the time between counts of about  $10/J$ . A plot (not shown) would reveal that by  $Jt = 100$  the oscillations in the quantum result have lost 2/3 of their amplitude, but that the quantum and classical approaches still give quite different results.

Finally, as soon as the atom numbers are monitored vigorously enough so that the classical evolution in the form of oscillations of the atoms between the two sides of the double-well trap can be resolved in detail in the scattered light (figures 3(b) and (d)), the quantum and classical descriptions give materially the same results. In round numbers, the time scale for classical evolution in our examples is  $1/J$  and the scattering rate to both detectors is  $N^2\Gamma/4$ . The watershed criterion is therefore  $N^2\Gamma/4J \sim 1$ , exceeded by a factor of about 10 in both graphs 3(b) and (d). Moreover, and unlike in the usual case with the ordinary TWA, in our



amended version of the TWA the quantum–classical agreement seems to persist for a long time, possibly as long as our numerical computations remain reliable.

The emergent classicality is a consequence of measurement back-action suppressing the subtle interference effects associated with quantum revivals that are inherent to the full quantum description [3]. Remarkably, this occurs even deep in the domain of quantum mechanics with only ten atoms, representing a far more dramatic transition to classicality than, e.g., measurement-induced relative phase between condensates [5–7] or nonlinear instabilities [10, 12, 13, 39]. The classical description at this level of accuracy is only possible because we have included classical phase diffusion to account for quantum mechanical measurement back-action. To wit, the only difference between the classical predictions in graphs 3(c) and (d) is the amount of phase diffusion, yet they are clearly different.

The main technical achievement of this paper is that we have introduced a theoretical method for a continuously monitored quantum system that handles both quantum fluctuations and measurement back-action classically. Now, the computational resources required to solve the time evolution from quantum mechanics tend to increase very rapidly with the system size. We, therefore, hope to inspire practical algorithms to analyze time evolution in situations where quantum effects are small but not negligible.

## Acknowledgments

This work was partially supported by the NSF (grant no. PHY-0967644), the Leverhulme Trust and the EPSRC.

## References

- [1] Joos E and Zeh H D 1985 The emergence of classical properties through interaction with the environment *Z. Phys. B* **59** 223–43
- [2] Zurek W H 2003 Decoherence, einselection, and the quantum origins of the classical *Rev. Mod. Phys.* **75** 715–75
- [3] Walls D F and Milburn G J 1985 Effect of dissipation on quantum coherence *Phys. Rev. A* **31** 2403–8
- [4] Leggett A J and Sols F 1991 On the concept of spontaneously broken gauge symmetry in condensed matter physics *Found. Phys.* **21** 353–64
- [5] Javanainen J and Yoo S M 1996 Quantum phase of a Bose–Einstein condensate with an arbitrary number of atoms *Phys. Rev. Lett.* **76** 161–4
- [6] Castin Y and Dalibard J 1997 Relative phase of two Bose–Einstein condensates *Phys. Rev. A* **55** 4330–7
- [7] Cirac J I, Gardiner C W, Naraschewski M and Zoller P 1996 Continuous observation of interference fringes from Bose condensates *Phys. Rev. A* **54** R3714–7
- [8] Spiller T P and Ralph J F 1994 The emergence of chaos in an open quantum system *Phys. Lett. A* **194** A235–40
- [9] Brun T A, Gisin N, O’Mahony P F and Rigo M 1997 From quantum trajectories to classical orbits *Phys. Lett. A* **229** A267–72
- [10] Bhattacharya T, Habib S and Jacobs K 2000 Continuous quantum measurement and the emergence of classical chaos *Phys. Rev. Lett.* **85** 4852–5
- [11] Chaudhury S, Smith A, Anderson B E, Ghose S and Jessen P S 2009 Quantum signatures of chaos in a kicked top *Nature* **461** 768
- [12] Scott A J and Milburn G J 2001 Quantum nonlinear dynamics of continuously measured systems *Phys. Rev. A* **63** 042101

- [13] Javanainen J and Shrestha U 2008 Nonlinear phenomenology from quantum mechanics: soliton in a lattice *Phys. Rev. Lett.* **101** 170405
- [14] Ruostekoski J and Walls D F 1998 Bose–Einstein condensate in a double-well potential as an open quantum system *Phys. Rev. A* **58** R50–3
- [15] Milburn G J, Corney J, Wright E M and Walls D F 1997 Quantum dynamics of an atomic Bose–Einstein condensate in a double-well potential *Phys. Rev. A* **55** 4318–24
- [16] Javanainen J 1986 Oscillatory exchange of atoms between traps containing Bose condensates *Phys. Rev. Lett.* **57** 3164–6
- [17] Raghavan S, Smerzi A, Fantoni S and Shenoy S R 1999 Coherent oscillations between two weakly coupled Bose–Einstein condensates: Josephson effects,  $\pi$  oscillations, and macroscopic quantum self-trapping *Phys. Rev. A* **59** 620–33
- [18] Albiez M, Gati R, Fölling J, Hunsmann S, Cristiani M and Oberthaler M K 2005 Direct observation of tunneling and nonlinear self-trapping in a single bosonic Josephson junction *Phys. Rev. Lett.* **95** 010402
- [19] Levy S, Lahoud E, Shomroni I and Steinhauer J 2007 The a.c. and d.c. Josephson effects in a Bose–Einstein condensate *Nature* **449** 579–83
- [20] Zibold T, Nicklas E, Gross C and Oberthaler M K 2010 Classical bifurcation at the transition from Rabi to Josephson dynamics *Phys. Rev. Lett.* **105** 204101
- [21] Lindblad G 1976 On the generators of quantum dynamical semigroups *Commun. Math. Phys.* **48** 119
- [22] Gardiner C and Zoller P 2004 *Quantum Noise* 3rd edn (Berlin: Springer)
- [23] Drummond P D and Hardman A D 1993 Simulation of quantum effects in Raman-active waveguides *Europhys. Lett.* **21** 279
- [24] Walls D F and Milburn G J 1994 *Quantum Optics* 2nd edn (Berlin: Springer)
- [25] Narducci L M, Alton Coulter C and Bowden C M 1974 Exact diffusion equation for a model for superradiant emission *Phys. Rev. A* **9** 829–45
- [26] Trimborn F, Witthaut D and Korsch H J 2008 Exact number-conserving phase-space dynamics of the  $m$ -site Bose–Hubbard model *Phys. Rev. A* **77** 043631
- [27] Trimborn F, Witthaut D and Korsch H J 2009 Beyond mean-field dynamics of small Bose–Hubbard systems based on the number-conserving phase-space approach *Phys. Rev. A* **79** 013608
- [28] Hush M R, Carvalho A R R and Hope J J 2012 Number-phase Wigner representation for scalable stochastic simulations of controlled quantum systems *Phys. Rev. A* **85** 023607
- [29] Dalibard J, Castin Y and Mølmer K 1992 Wave-function approach to dissipative processes in quantum optics *Phys. Rev. Lett.* **68** 580–3
- [30] Dum R, Zoller P and Ritsch H 1992 Monte Carlo simulation of the atomic master equation for spontaneous emission *Phys. Rev. A* **45** 4879–87
- [31] Tian L and Carmichael H J 1992 Quantum trajectory simulations of two-state behavior in an optical cavity containing one atom *Phys. Rev. A* **46** R6801–4
- [32] Wiseman H M and Milburn G J 2010 *Quantum Measurement and Control* (Cambridge: Cambridge University Press)
- [33] Isella L and Ruostekoski J 2006 Quantum dynamics in splitting a harmonically trapped Bose–Einstein condensate by an optical lattice: truncated Wigner approximation *Phys. Rev. A* **74** 063625
- [34] Blakie P B, Bradley A S, Davis M J, Ballagh R J and Gardiner C W 2008 Dynamics and statistical mechanics of ultra-cold Bose gases using c-field techniques *Adv. Phys.* **57** 363–455
- [35] Polkovnikov A 2010 Phase space representation of quantum dynamics *Ann. Phys., NY* **325** 1790–825
- [36] Martin A D and Ruostekoski J 2010 Quantum and thermal effects of dark solitons in a one-dimensional Bose gas *Phys. Rev. Lett.* **104** 194102
- [37] Gardiner C 2009 *Stochastic Methods* 4th edn (Berlin: Springer)
- [38] Ruostekoski J and Isella L 2005 Dissipative quantum dynamics of bosonic atoms in a shallow 1d optical lattice *Phys. Rev. Lett.* **95** 110403
- [39] Javanainen J 2010 Nonlinearity from quantum mechanics: dynamically unstable Bose–Einstein condensate in a double-well trap *Phys. Rev. A* **81** 051602